


# Protecting fresh waterways in Aotearoa/NZ: The strong Public Health case

16 October 2019

Nick Wilson, Leah Grout, Mereana Wilson , Anja Mizdrak , Phil Shoemack , Michael Baker



**Protecting waterways has the benefits of: (1) protecting water from hazardous microbes; (2) minimising cancer risk and other problems from nitrates in water; (3) avoiding algal blooms that are hazardous to health; (4) protecting mahinga kai uses (cultural importance and food security); (5) facilitating safe recreational water use; (6) minimising flooding risks from silted up waterways; and (7) protecting renewable energy from waterway sediments. In this blog we briefly consider these issues and why health workers and agencies should now do submissions on protecting waterways to the Ministry for the Environment, as part of a current consultation process which ends on 31 October.**

## Introduction

The NZ Ministry for the Environment (MfE) has recently published a discussion document, titled “Action for healthy waterways” [1]. As part of developing a submission on this document, we briefly review in this blog the public health case for waterways protection.



## **1) Protecting water from hazardous microbial contamination**

Dairy cattle, and other livestock, can have a major adverse impact on water quality [2]. Key sources of water pollution from dairy farms include animal wastes, pharmaceutical residues (eg, antibiotics, hormones), fertilisers and pesticides used for growing feed crops, and sediment from eroded pastures [2]. Antibiotic resistant bacteria and their genes can also act as environmental contaminants [3-5]. These pollutants can enter waterways directly through runoff from farm buildings, spills or the failure of manure storage facilities, the deposition of faecal matter into streams, and transport through soil layers via drainage waters on farms. Contamination can also occur indirectly from surface runoff and overland flow from pastures or agricultural fields [2].

Cattle excrete a number of different zoonotic pathogens that can contaminate the environment and cause illness in humans [2, 6, 7]. Once these pathogens enter the farm environment, they can infect humans via a range of different pathways [6-8]. A number of different pathogens can disperse in the environment and survive in soil or water, and ingestion or recreational contact with contaminated water is an important risk factor for several zoonotic diseases [6, 7].

Declining water quality has recently become an important issue in NZ with increased public concern and political attention. Over the past 30 years, dairy farming has become increasingly intensive and over the same period NZ's waterways have shown a range of indicators of degraded water quality [9-12]. Levels of nutrients, sediments, and faecal microbes are frequently elevated in catchments dominated by urban or agricultural land use [9, 11, 12].

In terms of specific diseases, water contaminated with livestock faeces has resulted in the world's largest water-borne outbreak of campylobacter infection in Havelock North (see this

blog: [13]). The outbreak in 2016, resulted in an estimated 5500 cases, 45 hospitalisations, and 3 deaths by the time it was over [14]. Total economic costs were estimated at \$21 million [15]. Prior to that outbreak, campylobacter infection in NZ (sporadic cases and outbreaks) had also been regularly linked to contaminated water supplies (eg, reviewed in: [16]). Giardia infection can also be spread via water in NZ (eg, [17]). Associations have been found between rainfall and increased notifications of both giardiasis and cryptosporidiosis [18]. The authors of this study concluded that: “this finding suggests that improving water supply quality in New Zealand could reduce vulnerability to the impact of climate change on protozoan diseases” [18].

Contaminated freshwater also imposes costs in terms of the need for water treatment. Such treatment costs many millions of dollars per year and indeed Auckland Council is planning to spend \$7 billion over 10 years to upgrade water infrastructure [1].

## **2) Minimising cancer risk and other problems from nitrates in water supplies**

Nitrogen is an important pollutant from dairy farms to surface waters, groundwater, and marine waters [19]. Excessive nitrate is a direct threat to human health [2, 19]. High levels of nitrate in drinking water can lead to the development of methaemoglobinaemia in infants [2, 20]. Nitrate toxicity has also been linked to miscarriages in pregnant women and certain forms of cancer in adults [2, 21, 22]. Specifically, the long-term consumption of nitrate in drinking water has been positively associated with a higher risk for thyroid disease and the following cancers: non-Hodgkin’s lymphoma, stomach, colorectal, bladder, breast, and ovarian cancers [23-30].

The association of nitrates with colorectal cancer is of particular concern, partly because this cancer is now the second commonest cause of cancer deaths in NZ. A European study gave an estimate of 4% of colorectal cancers in the European Union that could be attributed to nitrate exposure [31]. Applied to NZ, this attributable fraction would suggest exposure to nitrates in drinking water could be causing approximately 120 cases of colorectal cancer a year (out of the 3000 total) and 50 deaths (out of 1200 total). Quantifying the level of nitrate exposure in NZ drinking water and estimating the associated health risk is a relatively urgent research priority for this country (for additional comment see: [32]).

## **3) Avoiding algal blooms that are hazardous to health**

Nutrient loading and eutrophication from agriculture can also lead to harmful algal blooms in freshwater ecosystems, such as rivers and lakes, as well as in marine ecosystems [21, 22]. Harmful algal blooms are a concern for humans and animals that ingest contaminated water, inhale aerosolised toxins, or consume contaminated fish or shellfish [21, 22, 33]. For example, a red tide from a novel marine toxin in Wellington Harbour in 1998 was extremely toxic to fish and marine invertebrates and also caused respiratory distress in harbour bystanders [34]. Algal blooms can also be a problem in terms of adverse impact on tourism and real estate values (at least from the North American experience [35]).

## **4) Protecting mahinga kai uses - cultural importance and food security**

Streams, rivers and lakes provide habitats, materials and food sources to support mahinga

kai (the collection of wild food and food sources) for species such as: tuna-eels, waikoura-freshwater crayfish, kakahi-freshwater mussels, inanga-whitebait, watercress and puha. These traditional practices and food sources are of critical importance to Māori cultural wellbeing, food security and potential economic benefits (such as the commercial eel industry) which are also valued and practiced by other New Zealanders. It has been reported that a “number of hapū/iwi have already identified mahinga kai values and attributes in iwi management plans, regional planning documents, and kaupapa Māori assessment frameworks” [1, 36]. Furthermore, many iwi and hapu groups have employed a range of methods and approaches to monitor individual taonga species that are of critical importance to them and their tribal interest areas. But research shows these valuable cultural food resources are damaged by excessive nutrients, toxins and silt from soil erosion and in waterways [37]. In addition, kai moana resources around river mouths (fish and shellfish) are also harmed by waterway pollution [1] (though we recognise that this coastal issue is outside of the MfE Document on proposed freshwater reforms).

In terms of food security, one NZ study [38] modelled a scenario where foods were gathered and gifted and this was found to reduce the cost of a healthy diet in New Zealand by 10%. Therefore protection of mahinga kai and kai moana resources are important for cultural wellbeing, nutrition and *food security*.

## **5) Facilitating safe recreational water use**

New Zealanders enjoy swimming, kayaking, rowing, boating and fishing in the country’s waterways. For example, 8% of NZ adults report participating in sports and recreation activities on, in, and beside lakes, rivers, and streams in the past week [39]. These pursuits have benefits in terms of physical activity as well as mental health. Yet all these activities can be threatened via microbiological contamination of waterways (with users getting enteric infections), degradation from excess nutrients, and problems such as algal blooms (see above).

Furthermore, the recreational activities also contribute to tourist revenue, with tourists coming for trout fishing, kayaking and other water-based activities. The Taupo fishery alone has been estimated as worth about \$70 million annually, and Fish & Game reports the total value of recreational fisheries for the country is probably at least \$250 million [1]. The health aspect here is that this tourism supports employment and incomes – which are important contributors to health and wellbeing.

## **6) Minimising flooding risks from silted-up waterways**

NZ is particularly vulnerable to flooding owing to its geography (steep hill country) and building of houses on flood plains. As a result flooding impacts can be very high eg, insurance costs from the 2004 Manawatu floods were \$112 million [40], but the true economic cost was probably several times this amount. Also floods in NZ can harm both mental health and social capital – as per the Manawatu floods in 2004 [41]. Furthermore, flooding risks might be increasing with heavy rainfalls associated with climate change (though this may vary by region around the country). To minimise the risk of floods it is important that soil erosion from agricultural land and waterway silting is minimised. This again highlights the value of returning erosion-prone high country land to bush/forests and to fencing off and revegetating strips bordering streams and rivers (ie, riparian fencing and planting). Enhanced pest control (eg, of feral deer, pigs and goats) will also protect forest biomass and so potentially reduce run-off and flooding risks [42].



## 7) Protecting renewable energy from waterway sediments

Finally, climate change is a critical health issue and a key strategy for NZ to meet its international commitments to reduce greenhouse gas emissions is to generate electricity from renewable sources – including hydro dams. Yet sediments can reduce dam water storage capacity and accelerate wear on turbines. This is currently mainly a problem for smaller hydro schemes in NZ where the annual sediment yield exceeds 2000 t/km<sup>2</sup> [43]. But in the long term, sediment flow will also continue to reduce the storage capacity of larger hydro schemes in NZ and even now there are turbine replacement costs due to sediment (reported as occurring every 5 years at a cost of approximately \$1.3 million [1]).

### What health workers and agencies need to do now

Given all these health-related issues associated with waterways, there is a strong public health case for further action to protect them. Therefore we encourage submissions on the Ministry for the Environment's current discussion document [1] (due by 31 October 2019). In particular, we encourage arguments for very strong action to protect waterways and to build appreciation for the health benefits, cultural benefits and cost-savings that can arise from such actions. In a future blog we plan to comment on more specific aspects of MfE's Discussion Document.

### Summary

In summary the protection of waterways in Aotearoa/NZ is undoubtedly important from the perspectives of public health and cultural wellbeing. Additional costs to the agricultural sector for waterways protection need to be balanced with potential cost impacts elsewhere in terms of enteric disease treatment costs, water treatment costs, loss of wild food sources, tourism losses (eg, from impacts on trout fisheries), and the costs to the hydropower schemes from silt in waterways.

### References

1. Ministry for the Environment. Action for healthy waterways – A discussion document on national direction for our essential freshwater. Wellington: Ministry for the Environment, 2019.  
<https://www.mfe.govt.nz/sites/default/files/media/Fresh%20water/action-for-healthy-waterways.pdf>.
2. FAO, Steinfeld H, Gerber P, Wassenaar T, Castel V, Rosales M, de Haan C: Livestock's long shadow: environmental issues and options. Rome, Italy: Food and Agriculture Organization of the United Nations (FAO); 2006.
3. Tripathi V, Tripathi P: Antibiotic Resistance Genes: An Emerging Environmental Pollutant. In: Kesari K (eds). *Perspectives in Environmental Toxicology*. Springer; 2017: 183-201.
4. Aitken SL, Dilworth TJ, Heil EL, Nailor MD. Agricultural Applications for Antimicrobials. A Danger to Human Health: An Official Position Statement of the Society of Infectious Diseases Pharmacists. *Pharmacotherapy: The Journal of Human Pharmacology and Drug Therapy*. 2016;36(4):422-432.
5. Oliver SP, Murinda SE, Jayarao BM. Impact of antibiotic use in adult dairy cows on antimicrobial resistance of veterinary and human pathogens: a comprehensive review. *Foodborne Pathogens & Disease*. 2011;8(3):337-355.
6. Cavarani S. Cattle industry and zoonotic risk. *Veterinary Research Communications*.

2008;32 Suppl 1:S19-24.

7. McDaniel CJ, Cardwell DM, Moeller RB, Gray GC. Humans and Cattle: A Review of Bovine Zoonoses. *Vector Borne and Zoonotic Diseases*. 2014;14(1):1-19.
8. Toth JD, Aceto HW, Rankin SC, Dou Z. Short communication: Survey of animal-borne pathogens in the farm environment of 13 dairy operations. *Journal of Dairy Science*. 2013;96(9):5756-5761.
9. Scarsbrook MR, Melland AR. Dairying and water-quality issues in Australia and New Zealand. *Animal Production Science*. 2015;55(7):856-868.
10. Ballantine D, Davies-Colley R. Water quality trends in New Zealand rivers: 1989-2009. *Environmental Monitoring and Assessment*. 2014;186(3):1939-1950.
11. Larned S, Scarsbrook M, Snelder T, Norton N, Biggs B. Water quality in low-elevation streams and rivers of New Zealand: recent state and trends in contrasting land-cover classes. *New Zealand Journal of Marine and Freshwater Research*. 2004;38(2):347-366.
12. Ministry for the Environment, Statistics NZ. *New Zealand's Environmental Reporting Series: Our fresh water 2017*. Ministry for the Environment, Statistics NZ. 2017.
13. Baker M, Wilson N, Woodward A. The Havelock North drinking water inquiry: A wake-up call to rebuild public health in New Zealand. *Public Health Expert [Blog]*. 2017 (20 December).  
<https://blogs.otago.ac.nz/pubhealthexpert/2017/12/20/the-havelock-north-drinking-water-inquiry-a-wake-up-call-to-rebuild-public-health-in-new-zealand/>.
14. New Zealand Government. Government Inquiry into Havelock North Drinking Water: Report of the Havelock North Drinking Water Inquiry: Stage 1. Auckland; 2017, May.
15. Moore D, Drew R, Davies P, Rippon R. The Economic Costs of the Havelock North August 2016 Waterborne Disease Outbreak.. Wellington: Sapere Research Group; 2017, August.
16. Wilson N. A Systematic Review of the Aetiology of Human Campylobacteriosis in New Zealand. Wellington: NZ Food Safety Authority, 2005.  
<https://www.mpi.govt.nz/dmsdocument/22330/direct>.
17. Hoque ME, Hope VT, Kjellstrom T, Scragg R, Lay-Yee R. Risk of giardiasis in Aucklanders: a case-control study. *International Journal of Infectious Diseases*. 2002;6(3):191-197.
18. Britton E, Hales S, Venugopal K, Baker MG. The impact of climate variability and change on cryptosporidiosis and giardiasis rates in New Zealand. *Journal of Water and Health*. 2010;8(3):561-571.
19. OECD: The Dairy Sector. In: *Agriculture, Trade and the Environment*. Paris, France: Organisation for Economic Co-operation and Development; 2004.
20. Gupta SK, Gupta RC, Seth AK, Gupta AB, Bassin JK, Gupta A. Adaptation of cytochrome-b 5 reductase activity and methaemoglobinaemia in areas with a high nitrate concentration in drinking-water. *Bulletin of the World Health Organization*. 1999;77(9):749-753.
21. Johnson PTJ, Townsend AR, Cleveland CC, Glibert PM, Howarth RW, McKenzie VJ, Rejmankova E, Ward MH. Linking environmental nutrient enrichment and disease emergence in humans and wildlife. *Ecological Applications*. 2010;20(1):16-29.
22. Townsend AR, Howarth RW, Bazzaz FA, Booth MS, Cleveland CC, Collinge SK, Dobson AP, Epstein PR, Holland EA, Keeney DR *et al*: Human health effects of a changing global nitrogen cycle. *Frontiers in Ecology of the Environment* 2003;1:240-246.
23. Jones RR, Weyer PJ, Dellavalle CT, Inoue-Choi M, Anderson KE, Cantor KP, Krasner S, Robien K, Freeman LEB, Silverman DT *et al*. Nitrate from Drinking Water and Diet and Bladder Cancer Among Postmenopausal Women in Iowa. *Environmental Health Perspectives*. 2016;124(11):1751.

24. Inoue-Choi M, Anderson K, Cerhan, Jr., Weyer P, Ward M. Dietary intake of nitrate and nitrite, nitrate in drinking water, and ovarian cancer risk among postmenopausal women in Iowa. *Cancer Research*. 2013;73(8).
25. Gulis G, Czompolyova M, Cerhan JR. An Ecologic Study of Nitrate in Municipal Drinking Water and Cancer Incidence in Trnava District, Slovakia. *Environmental Research*. 2002;88(3):182-187.
26. Fachiroh J, Gravitiani E, Husodo AH. Nitrate in drinking water and risk of colorectal cancer in Yogyakarta, Indonesia. *Journal of Toxicology and Environmental Health Part A*. 2017;80(2):120-128.
27. Espejo-Herrera N, Gràcia-Lavedan E, Boldo E, Aragonés N, Pérez-Gómez B, Pollán M, Molina AJ, Fernández T, Martín V, La Vecchia C *et al*. Colorectal cancer risk and nitrate exposure through drinking water and diet. *International Journal of Cancer*. 2016;139(2):334-346.
28. Weyer PJ, Cerhan JR, Kross BC, Hallberg GR, Kantamneni J, Breuer G, Jones MP, Zheng W, Lynch CF. Municipal Drinking Water Nitrate Level and Cancer Risk in Older Women: The Iowa Women's Health Study. *Epidemiology*. 2001;12(3):327-338.
29. Schullehner J, Hansen B, Thygesen M, Pedersen CB, Sigsgaard T. Nitrate in drinking water and colorectal cancer risk: A nationwide population-based cohort study. *International Journal of Cancer*. 2018;143(1):73-79.
30. Ward MH, Jones RR, Brender JD, de Kok TM, Weyer PJ, Nolan BT, Villanueva CM, van Breda SG. Drinking Water Nitrate and Human Health: An Updated Review. *International Journal of Environmental Research and Public Health*. 2018;15(7).
31. van Grinsven HJ, Rabl A, de Kok TM. Estimation of incidence and social cost of colon cancer due to nitrate in drinking water in the EU: a tentative cost-benefit assessment. *Environmental Health*. 2010;9:58.
32. Joy M, Baker M. Drinking water study raises health concerns for New Zealanders. *The Conversation* 2019;(25 January).  
<https://theconversation.com/drinking-water-study-raises-health-concerns-for-new-zealanders-108510>.
33. Ministry of Health: Guidelines for Drinking-water Quality Management for New Zealand Inc., 5th edn. Wellington, New Zealand: Ministry of Health; 2017.
34. Hamamoto Y, Tachibana K, Holland PT, Shi F, Beuzenberg V, Itoh Y, Satake M. Brevisulcinal-F: a polycyclic ether toxin associated with massive fish-kills in New Zealand. *Journal of the American Chemical Society*. 2012;134(10):4963-4968.
35. Carmichael WW, Boyer GL. Health impacts from cyanobacteria harmful algae blooms: Implications for the North American Great Lakes. *Harmful Algae*. 2016;54:194-212.
36. Rainforth HJ, Harmsworth GR. Kaupapa Māori Freshwater Assessments: A summary of iwi and hapū-based tools, frameworks and methods for assessing freshwater environments. Perception Planning Ltd, 2019. pp115.
37. Stewart M, Tipa G, Williams E, Home M, Olsen G, Hickey C. Impacts of bioaccumulative contaminants in the Te Waihora catchment on Mahinga Kai gatherers: data report and risk assessment. NIWA Report HAM2014-012, National Institute of Water and Atmospheric Research, Hamilton, New Zealand, 2014.
38. Mackay S, Buch T, Vandevijvere S, Goodwin R, Korohina E, Funaki-Tahifote M, Lee A, Swinburn B. Cost and Affordability of Diets Modelled on Current Eating Patterns and on Dietary Guidelines, for New Zealand Total Population, Maori and Pacific Households. *International Journal of Environmental Research and Public Health*. 2018;15(6).
39. Sport New Zealand. Active NZ 2017 Participation Report. Wellington: Sport New Zealand, 2018.
40. Grocott M. Suddenly it was not practice. *Manawatu Standard* 2014 (15 February).  
<http://i.stuff.co.nz/manawatu-standard/9726105/Suddenly-it-was-not-practice>.

41. Smith W, Davies-Colley C, Mackay A, Bankoff G. Social impact of the 2004 Manawatu floods and the 'hollowing out' of rural New Zealand. *Disasters*. 2011;35(3):540-553.
42. Wilson N, Blaschke P, Thomson G, Nghiem N, Horrocks J. Public health aspects of feral deer, goats and pigs in New Zealand: A review to inform eradication decisions. *New Zealand Geographer*. 2015;71:177-188.
43. Jowett I. Sedimentation in New Zealand hydroelectric schemes. *Water International*. 1984;9:172-176.

Public Health Expert Briefing (ISSN 2816-1203)

---

**Source URL:**

<https://www.phcc.org.nz/briefing/protecting-fresh-waterways-aotearoanz-strong-public-health-case>